ELSEVIER

Journal of Organometallic Chemistry 543 (1997) 271

Journal of Organometallic Chemistry

Author Index of Volume 543

Ajulu, F.A., 245
Almansour, A.I., 83

Baumeister, U., 117
Beck, W., 39, 47
Beckmann, J., 229
Beer, P.D., 259
Berke, H., 171
Blake, A.J., 217
Bochmann, M., 209
Brauer, D., 103

Calderazzo, F., 201
Calhorda, M.J., 93
Canovese, L., 145
Caro, B., 87
Cheng, J., 227
Clark, A.M., 111
Contel, M., 71
Crociani, B., 145
Cuenca, T., 209
Cui, X.L., 63

Dash, A.K., 135
Drew, M.G.B., 259

Englert, U., 201

Fu, H., 227
Furukawa, N., 77

Galakhov, M., 209
Garrido, J., 71
Gauss, C., 171
Ghose, S., 189
Gibbons, M.N., 217
Gielen, R., 103
Gupta, B.D., 125

Hassler, K., 103, 117
Heck, J., 153
Herrmann, R., 93
Holle, S., 255
Holloway, C.E., 1
Hor, T.S.A., 179
Horn, E., 77
Hossain, Md.M., 135, 189
Hou, J., 227
Huang, X., 165
Hüffer, S., 47
Hussain, Z.I., 237

James, S.L., 233
Jänicke, M., 171
Jiang, C., 179
Jiménez, G., 209
Jiménez, J., 71
Jolly, P.W., 255
Jones, P.G., 71
Jurkschat, K., 229

Katsukawa, Y., 241
Katzenbeisser, U., 103
Kiang, F.-M., 249
Kirin, V.P., 57
Kölle, U., 201
Kollegger, G.M., 103
Koller, A., 171
Körnich, J., 153
Krüger, C., 103
Ku, R.-Z., 249
Kuhnigk, J., 255

Laguna, A., 71
Laguna, M., 71
Leadbeater, N.E., 251
Lewis, J., 233, 251
Liable-Sands, L.M., 135
Liu, C.-Y., 249
Liu, L.-K., 179
Liu, S.-T., 249
Liu, Y.H., 63
Lopes, P.E.M., 93
Lu, S., 227

Mahon, M.F., 189
Maksakov, V.A., 57
Mao, X.A., 63
Mathey, F., 245
Mathur, P., 135, 189
McInnes, E.J.L., 237
Melnik, M., 1
Milke, J., 39
Muto, H., 241

Nixon, J.F., 245

Onaka, S., 241

Pampaloni, G., 201
Podberezskaya, N.V., 57
Polborn, K., 47
Prat, V., 87

Qanungo, K., 125

Raithby, P.R., 233, 251
Rheingold, A.L., 135
Rickard, C.E.F., 111
Robin-Le Guen, F., 87
Roper, W.R., 111
Royo, E., 209
Royo, P., 209

Sato, S., 77
Satyanarayana, C.V.V., 135, 189
Schenzel, K., 117
Schier, A., 93
Schollmeyer, D., 229
Schürmann, M., 229
Semyannikov, P.P., 57
Sénéchal-Tocquer, M.-C., 87
Smith, D.K., 259
Sowerby, D.B., 217
Sünkel, K., 39

Takagi, S., 241
Tripepi, G., 201

Ueminami, T., 77
Uguagliati, P., 145

Vaissermann, J., 87
Virovets, A.V., 57
Visentin, F., 145

Wang, H., 227
Wen, Y.-S., 179
Whiteley, M.W., 237
Wright, L.J., 111
Wu, Y.J., 63

Yan, Y.K., 179
Yang, D.Y., 165
Yap, G.P.A., 135
Younus, M., 233
Yuan, H.Z., 63

Zhou, H., 227
Zink, R., 117

Elsevier Science S.A.

ELSEVIER

Journal of Organometallic Chemistry 543 (1997) 273–276

Journal of Organometallic Chemistry

Subject Index of Volume 543

Acetate
The preparation and reactions of $(\eta^3$-allyl$)_2$Ru[Pr$^i_2$P(CH$_2$)$_n$PPr$^i_2$] complexes (S. Holle, P.W. Jolly and J. Kuhnigk), 255

Acetylenic $Co_2(CO)_6$ fragment
Synthesis, structure and reactivity of acetylenic $Co_2(CO)_6$ pyrylium salts. Electronic influence of the $Co_2(CO)_6$ acetylenic fragment (B. Caro, F. Robin-Le Guen, M.-C. Sénéchal-Tocquer, V. Prat and J. Vaissermann), 87

(Alkylseleno)haloethenes
Stereoselective synthesis of conjugated alkadienes via the palladium-catalyzed coupling reaction of (*Z*)- or (*E*)-alkenylboranes with (*Z*)- or (*E*)-2-halo-1-(alkylseleno)ethenes (D.Y. Yang and X. Huang), 165

Alkenylboranes
Stereoselective synthesis of conjugated alkadienes via the palladium-catalyzed coupling reaction of (*Z*)- or (*E*)-alkenylboranes with (*Z*)- or (*E*)-2-halo-1-(alkylseleno)ethenes (D.Y. Yang and X. Huang), 165

Alkynyl bridge
Synthesis and properties of the alkynyl-bridged heterobimetallics $[Mo(C{=}CFc)(Ph_2PCH_2CH_2PPh_2)(\eta\text{-}C_7H_7)]^z$ (Fc = ferrocenyl, $z = 0$ or $+1$) (Z.I. Hussain, M.W. Whiteley and E.J.L. McInnes), 237

Alkynyl complexes
Kohlenwasserstoffverbrückte Metallkomplexe, XXXV. Addition von *tert*. Butylethinyl–pentacarbonyl–wolframat an π-koordinierte Kohlenwasserstoffe von kationischen Komplexen (J. Milke, K. Sünkel and W. Beck), 39

Asymmetric
Asymmetric hydroesterification of styrene by $PdCl_2$–$CuCl_2$-chiral phosphine catalyst systems (H. Zhou, J. Hou, J. Cheng, S. Lu, H. Fu and H. Wang), 227

Bis(cyclopentadienyl)-bridged complexes
Kooperative Wirkung in π-Ligand-verbrückten Zweikernkomplexen XX. Eine neue Strategie zur Synthese von heterodinuklearen Komplexen mit einem verbrückenden Bis(cyclopentadienyl)-Liganden [2] (J. Heck und J. Körnich), 153

Book Review
Advances in Physical Organic Chemistry, D. Bethell (Ed.) (C. Eaborn), 263
Advances in transition metal coordination chemistry, Chi-Ming Che and Vivian W.W. Yam (M.R. Lappert), 264
Applied homogeneous catalysis with organometallic compounds, B. Cornils andW.A. Herrmann (Eds.) (M.F. Tappert), 264
Gmelin Handbook of Inorganic and Organometallic Chemistry. Sn. Organotin Compounds. Part 24. Dinuclear Compounds, Springer-Verlag, Berlin (P.A. Chaloner), 263
Palladium Reagents and Catalysis; Innovations in Organic Synthesis, Jiro Tsuji, John Wiley and Sons (Eds.) (P.A. Chaloner), 268
Perspectives on Bioinorganic Chemistry, R.W. Hay, J.R. Dilworth, and K.B. Nolan (Eds.) (G.J. Leigh), 265
Synthetic Methods of Organometallic and Inorganic Chemistry, F.T. Edelmann (Ed.) (J.D. Smith), 267
The Molecular Chemistry of the Transition Elements, F. Mathey and A. Sevin (Eds.) (G.J. Leigh), 265
Topics in Current Chemistry Volume 179 Organolanthanoid Chemistry: Synthesis, Structure, Catalysis. Springer, Berlin (G.A. Lawless), 266

Carbonyl
Kohlenwasserstoffverbrückte Metallkomplexe, XXXVI. Nucleophile Addition von Carbonylmetallaten an koordinierte Ethen-, Benzol-, Cyclooctadienyl- und Cyclooctatrienyl-Liganden von kationischen Ruthenium-Komplexen (S. Hüffer, K. Polborn and W. Beck), 47
Synthesis and spectroscopic characterization of $(CO)_6Fe_2(\mu\text{-}EC(H){=}C(2\text{-}Th)E')$ (E,E' = S, Se, Te; Th = C_4H_3S). Structural characterization of $(CO)_6Fe_2(\mu\text{-}SeC(H){=}C(2\text{-}Th)Se)$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana, A.L. Rheingold, G.P.A. Yap and L.M. Liable-Sands), 135

Carbonyl-hydroxo-alkoxo-complex
Amine-oxide-mediated reactions of $Re_2(CO)_{10}$ with phenol and aliphatic alcohols: The formation of $Re_3(CO)_{15}(\mu\text{-H})$ and a hydroxo-methoxo trirhenium aggregate $[Me_3NH]^+[Re_3(CO)_9(\mu\text{-}OH)_2(\mu\text{-}OMe)(\mu_3\text{-}OMe)]^-$ (C. Jiang, Y.-S. Wen, L.-K. Liu, T.S.A. Hor and Y.K. Yan), 179

Catalysis
Stereoselective synthesis of conjugated alkadienes via the palladium-catalyzed coupling reaction of (*Z*)- or (*E*)-alkenylboranes with (*Z*)- or (*E*)-2-halo-1-(alkylseleno)ethenes (D.Y. Yang and X. Huang), 165

Cationic dinuclear derivatives
Dinuclear cationic zirconium complexes with the fulvalene ligand. Synthesis and reactivity (T. Cuenca, M. Galakhov, G. Jiménez, E. Royo, P. Royo and M. Bochmann), 209

Clusters
Thermolysis of $Fe_3(CO)_9(\mu_3\text{-}Se)(\mu_3\text{-}E)$ (E = S, Te) with $Cp_2Mo_2(CO)_6$ and formation of new mixed-chalcogenide, mixed Fe–Mo carbonyl clusters. Crystal structures of $Cp_2Mo_2Fe_2(CO)_6(\mu_3\text{-}Se)_2(\mu_4\text{-}Te)$, $Cp_2Mo_2Fe_2(CO)_7(\mu_3\text{-}Se)(\mu_3\text{-}Te)$ and $Cp_2Mo_2Fe_2(CO)_6(\mu_3\text{-}S)(\mu_3\text{-}Se)(\mu_4\text{-}Se)$ (P. Mathur, S. Ghose, Md.M. Hossain, C.V.V. Satyanarayana and M.F. Mahon), 189

Cobalt cluster
The unexpected synthesis of a tricobalt cluster with ferrocenyl groups: crystal structure and cyclic voltammetry of $Co_3(CO)_7(\mu_3\text{-}PFc)(\mu_2\text{-}PFc_2)$ (Fc = $(C_5H_5)Fe(C_5H_4)$) (S. Onaka, H. Muto, Y. Katsukawa and S. Takagi), 241

Coupling
Stereoselective synthesis of conjugated alkadienes via the palladium-catalyzed coupling reaction of (*Z*)- or (*E*)-alkenylboranes with

Elsevier Science S.A.

(*Z*)- or (*E*)-2-halo-1-(alkylseleno)ethenes (D.Y. Yang and X. Huang), 165

Crystal structure
Cyclometallation of ferrocenylimines. III. Regioselectivity in Hg(II) cyclometallated complexes (Y.J. Wu, X.L. Cui, Y.H. Liu, H.Z. Yuan and X.A. Mao), 63
Isolation and X-ray structure determination of a new tellurane *Te*-oxide dimer [12-Te-6(C4O2)]$_2$ (λ^6-tellane) (S. Sato, T. Ueminami, E. Horn and N. Furukawa), 77

Cyclic voltammetry
The unexpected synthesis of a tricobalt cluster with ferrocenyl groups: crystal structure and cyclic voltammetry of $Co_3(CO)_7(\mu_3\text{-}PFc)(\mu_2\text{-}PFc_2)$ (Fc = $(C_5H_5)Fe(C_5H_4)$) (S. Onaka, H. Muto, Y. Katsukawa and S. Takagi), 241

Cycloheptatrienyl
Synthesis and properties of the alkynyl-bridged heterobimetallics $[Mo(C{\equiv}CFc)(Ph_2PCH_2CH_2PPh_2)(\eta\text{-}C_7H_7)]^z$ (Fc = ferrocenyl, $z = 0$ or $+1$) (Z.I. Hussain, M.W. Whiteley and E.J.L. McInnes), 237

Cyclopentadienyl derivatives
Dinuclear cationic zirconium complexes with the fulvalene ligand. Synthesis and reactivity (T. Cuenca, M. Galakhov, G. Jiménez, E. Royo, P. Royo and M. Bochmann), 209

Decarbonylation
Amine-oxide-mediated reactions of $Re_2(CO)_{10}$ with phenol and aliphatic alcohols: The formation of $Re_3(CO)_{14}(\mu\text{-}H)$ and a hydroxo-methoxo trirhenium aggregate $[Me_3NH]^+[Re_3(CO)_9(\mu\text{-}OH)_2(\mu\text{-}OMe)(\mu_3\text{-}OMe)]^-$ (C. Jiang, Y.-S. Wen, L.-K. Liu, T.S.A. Hor and Y.K. Yan), 179

Dienyl selenides
Stereoselective synthesis of conjugated alkadienes via the palladium-catalyzed coupling reaction of (*Z*)- or (*E*)-alkenylboranes with (*Z*)- or (*E*)-2-halo-1-(alkylseleno)ethenes (D.Y. Yang and X. Huang), 165

Dirhenium decacarbonyl
Amine-oxide-mediated reactions of $Re_2(CO)_{10}$ with phenol and aliphatic alcohols: The formation of $Re_3(CO)_{14}(\mu\text{-}H)$ and a hydroxo-methoxo trirhenium aggregate $[Me_3NH]^+[Re_3(CO)_9(\mu\text{-}OH)_2(\mu\text{-}OMe)(\mu_3\text{-}OMe)]^-$ (C. Jiang, Y.-S. Wen, L.-K. Liu, T.S.A. Hor and Y.K. Yan), 179

Electrochemistry
Synthesis and properties of the alkynyl-bridged heterobimetallics $[Mo(C{\equiv}CFc)(Ph_2PCH_2CH_2PPh_2)(\eta\text{-}C_7H_7)]^z$ (Fc = ferrocenyl, $z = 0$ or $+1$) (Z.I. Hussain, M.W. Whiteley and E.J.L. McInnes), 237

Esr spectroscopy
Synthesis and properties of the alkynyl-bridged heterobimetallics $[Mo(C{\equiv}CFc)(Ph_2PCH_2CH_2PPh_2)(\eta\text{-}C_7H_7)]^z$ (Fc = ferrocenyl, $z = 0$ or $+1$) (Z.I. Hussain, M.W. Whiteley and E.J.L. McInnes), 237

Ferrocene
Cyclometallation of ferrocenylimines. III. Regioselectivity in Hg(II) cyclometallated complexes (Y.J. Wu, X.L. Cui, Y.H. Liu, H.Z. Yuan and X.A. Mao), 63

Ferrocenyl groups
The unexpected synthesis of a tricobalt cluster with ferrocenyl groups: crystal structure and cyclic voltammetry of $Co_3(CO)_7(\mu_3\text{-}PFc)(\mu_2\text{-}PFc_2)$ (Fc = $(C_5H_5)Fe(C_5H_4)$) (S. Onaka, H. Muto, Y. Katsukawa and S. Takagi), 241

Group VI metals
Oxidative cleavage of metal carbene complexes by iodine (S.-T. Liu, R.-Z. Ku, C.-Y. Liu and F.-M. Kiang), 249

η^3-allyl
The preparation and reactions of $(\eta^3\text{-allyl})_2Ru[Pr^i_2P(CH_2)_nPPr^i_2]$ complexes (S. Holle, P.W. Jolly and J. Kuhnigk), 255

Heterobimetallic
Synthesis and properties of the alkynyl-bridged heterobimetallics $[Mo(C{\equiv}CFc)(Ph_2PCH_2CH_2PPh_2)(\eta\text{-}C_7H_7)]^z$ (Fc = ferrocenyl, $z = 0$ or $+1$) (Z.I. Hussain, M.W. Whiteley and E.J.L. McInnes), 237

Heterodinuclear complexes
Kooperative Wirkung in π-Ligand-verbrückten Zweikernkomplexen XX. Eine neue Strategie zur Synthese von heterodinuklearen Komplexen mit einem verbrückenden Bis(cyclopentadienyl)-Liganden [2] (J. Heck und J. Körnich), 153

Heterometallic clusters
Synthesis of trinuclear mixed-metal clusters $Os_2Fe(CO)_{10}L_2$ (L = OH, I, CO). The crystal structures of $Os_2Fe(CO)_{10}(\mu\text{-}OH)_2$ and $Os_2Fe(CO)_{12} \cdot 0.41Os_3(CO)_{12}$ (V.A. Maksakov, V.P. Kirin, A.V. Virovets, N.V. Podberezskaya and P.P. Semyannikov), 57

Hexacoordinate
Isolation and X-ray structure determination of a new tellurane *Te*-oxide dimer [12-Te-6(C4O2)]$_2$ (λ^6-tellane) (S. Sato, T. Ueminami, E. Horn and N. Furukawa), 77

η^5-1-hexadienyl
The preparation and reactions of $(\eta^3\text{-allyl})_2Ru[Pr^i_2P(CH_2)_nPPr^i_2]$ complexes (S. Holle, P.W. Jolly and J. Kuhnigk), 255

Homodinuclear complexes
Kooperative Wirkung in π-Ligand-verbrückten Zweikernkomplexen XX. Eine neue Strategie zur Synthese von heterodinuklearen Komplexen mit einem verbrückenden Bis(cyclopentadienyl)-Liganden [2] (J. Heck und J. Körnich), 153

Hydride
Amine-oxide-mediated reactions of $Re_2(CO)_{10}$ with phenol and aliphatic alcohols: The formation of $Re_3(CO)_{14}(\mu\text{-}H)$ and a hydroxo-methoxo trirhenium aggregate $[Me_3NH]^+[Re_3(CO)_9(\mu\text{-}OH)_2(\mu\text{-}OMe)(\mu_3\text{-}OMe)]^-$ (C. Jiang, Y.-S. Wen, L.-K. Liu, T.S.A. Hor and Y.K. Yan), 179

Hydrocarbon bridges
Kohlenwasserstoffverbrückte Metallkomplexe, XXXVI. Nucleophile Addition von Carbonylmetallaten an koordinierte Ethen-, Benzol-, Cyclooctadienyl- und Cyclooctatrienyl-Liganden von kationischen Ruthenium-Komplexen (S. Hüffer, K. Polborn and W. Beck), 47

Hydroesterification
Asymmetric hydroesterification of styrene by $PdCl_2$–$CuCl_2$-chiral phosphine catalyst systems (H. Zhou, J. Hou, J. Cheng, S. Lu, H. Fu and H. Wang), 227

Imidazolidin-2-ylidinium iodides
Oxidative cleavage of metal carbene complexes by iodine (S.-T. Liu, R.-Z. Ku, C.-Y. Liu and F.-M. Kiang), 249

Iodine
Oxidative cleavage of metal carbene complexes by iodine (S.-T. Liu, R.-Z. Ku, C.-Y. Liu and F.-M. Kiang), 249

Iron
Kohlenwasserstoffverbrückte Metallkomplexe, XXXVI. Nucleophile Addition von Carbonylmetallaten an koordinierte Ethen-, Benzol-, Cyclooctadienyl- und Cyclooctatrienyl-Liganden von kationischen Ruthenium-Komplexen (S. Hüffer, K. Polborn and W. Beck), 47
Synthesis and spectroscopic characterization of $(CO)_6Fe_2(\mu\text{-}EC(H){=}C(2\text{-}Th)E')$ (E,E' = S, Se, Te; Th = C_4H_3S). Structural characterization of $(CO)_6Fe_2(\mu\text{-}SeC(H){=}C(2\text{-}Th)Se)$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana, A.L. Rheingold, G.P.A. Yap and L.M. Liable-Sands), 135
Synthesis of trinuclear mixed-metal clusters $Os_2Fe(CO)_{10}L_2$ (L = OH, I, CO). The crystal structures of $Os_2Fe(CO)_{10}(\mu\text{-}OH)_2$ and

$Os_2Fe(CO)_{12} \cdot 0.41Os_3(CO)_{12}$ (V.A. Maksakov, V.P. Kirin, A.V. Virovets, N.V. Podberezskaya and P.P. Semyannikov), 57

Thermolysis of $Fe_3(CO)_9(\mu_3\text{-}Se)(\mu_3\text{-}E)$ (E = S, Te) with $Cp_2Mo_2(CO)_6$ and formation of new mixed-chalcogenide, mixed Fe–Mo carbonyl clusters. Crystal structures of $Cp_2Mo_2Fe_2(CO)_6(\mu_3\text{-}Se)_2(\mu_4\text{-}Te)$, $Cp_2Mo_2Fe_2(CO)_7(\mu_3\text{-}Se)(\mu_3\text{-}Te)$ and $Cp_2Mo_2Fe_2(CO)_6(\mu_3\text{-}S)(\mu_3\text{-}Se)(\mu_4\text{-}Se)$ (P. Mathur, S. Ghose, Md.M. Hossain, C.V.V. Satyanarayana and M.F. Mahon), 189

Mechanisms

Comparison of the reactivities of $(Me_3Si)_3CSiMe_2I$ and $(Pr^iMe_2Si)_3CSiMe_2I$ (A.I. Almansour), 83

Mercury

Cyclometallation of ferrocenylimines. III. Regioselectivity in Hg(II) cyclometallated complexes (Y.J. Wu, X.L. Cui, Y.H. Liu, H.Z. Yuan and X.A. Mao), 63

Metal carbene complexes

Oxidative cleavage of metal carbene complexes by iodine (S.-T. Liu, R.-Z. Ku, C.-Y. Liu and F.-M. Kiang), 249

Methyl propiolate

Methyl propiolate coupling at the $Fe(CO)_2(PEt_3)_2$ fragment (M. Jänicke, C. Gauss, A. Koller and H. Berke), 171

Molybdenum

Synthesis and properties of the alkynyl-bridged heterobimetallics $[Mo(C{\equiv}CFc)(Ph_2PCH_2CH_2PPh_2)(\eta\text{-}C_7H_7)]^z$ (Fc = ferrocenyl, z = 0 or +1) (Z.I. Hussain, M.W. Whiteley and E.J.L. McInnes), 237

Thermolysis of $Fe_3(CO)_9(\mu_3\text{-}Se)(\mu_3\text{-}E)$ (E = S, Te) with $Cp_2Mo_2(CO)_6$ and formation of new mixed-chalcogenide, mixed Fe–Mo carbonyl clusters. Crystal structures of $Cp_2Mo_2Fe_2(CO)_6(\mu_3\text{-}Se)_2(\mu_4\text{-}Te)$, $Cp_2Mo_2Fe_2(CO)_7(\mu_3\text{-}Se)(\mu_3\text{-}Te)$ and $Cp_2Mo_2Fe_2(CO)_6(\mu_3\text{-}S)(\mu_3\text{-}Se)(\mu_4\text{-}Se)$ (P. Mathur, S. Ghose, Md.M. Hossain, C.V.V. Satyanarayana and M.F. Mahon), 189

Osmium

Kohlenwasserstoffverbrückte Metallkomplexe, XXXVI. Nucleophile Addition von Carbonylmetallaten an koordinierte Ethen-, Benzol-, Cyclooctadienyl- und Cyclooctatrienyl-Liganden von kationischen Ruthenium-Komplexen (S. Hüffer, K. Polborn and W. Beck), 47

Synthesis of trinuclear mixed-metal clusters $Os_2Fe(CO)_{10}L_2$ (L = OH, I, CO). The crystal structures of $Os_2Fe(CO)_{10}(\mu\text{-}OH)_2$ and $Os_2Fe(CO)_{12} \cdot 0.41Os_3(CO)_{12}$ (V.A. Maksakov, V.P. Kirin, A.V. Virovets, N.V. Podberezskaya and P.P. Semyannikov), 57

Osmium nitrosyl complexes

Osmium nitrosyl complexes with osmium–tin bonds. Crystal structure of $Os[Sn(p\text{-tolyl})_3](NO)(CO)_2(PPh_3)$ (A.M. Clark, C.E.F. Rickard, W.R. Roper and L.J. Wright), 111

Osmium-tin bonds

Osmium nitrosyl complexes with osmium–tin bonds. Crystal structure of $Os[Sn(p\text{-tolyl})_3](NO)(CO)_2(PPh_3)$ (A.M. Clark, C.E.F. Rickard, W.R. Roper and L.J. Wright), 111

Oxidative cleavage

Oxidative cleavage of metal carbene complexes by iodine (S.-T. Liu, R.-Z. Ku, C.-Y. Liu and F.-M. Kiang), 249

Palladium

Stereoselective synthesis of conjugated alkadienes via the palladium-catalyzed coupling reaction of (*Z*)- or (*E*)-alkenylboranes with (*Z*)- or (*E*)-2-halo-1-(alkylseleno)ethenes (D.Y. Yang and X. Huang), 165

$PdCl_2$–$CuCl_2$-chiral phosphine catalyst systems

Asymmetric hydroesterification of styrene by $PdCl_2$–$CuCl_2$-chiral phosphine catalyst systems (H. Zhou, J. Hou, J. Cheng, S. Lu, H. Fu and H. Wang), 227

Pertellurane

Isolation and X-ray structure determination of a new tellurane *Te*-oxide dimer $[12\text{-}Te\text{-}6(C4O2)]_2$ (λ^6-tellane) (S. Sato, T. Ueminami, E. Horn and N. Furukawa), 77

Polymer

On the reaction of $[Ph_2(OH)Si]_2O$ with t-Bu_2SnCl_2: synthesis and characterization of the first well defined polystannasiloxane $[(t\text{-}Bu_2SnO)(Ph_2SiO)_2]_n$ (J. Beckmann, K. Jurkschat, D. Schollmeyer and M. Schürmann), 229

Protolysis

Amine-oxide-mediated reactions of $Re_2(CO)_{10}$ with phenol and aliphatic alcohols: The formation of $Re_3(CO)_{14}(\mu\text{-}H)$ and a hydroxo-methoxo trirhenium aggregate $[Me_3NH]^+[Re_3(CO)_9(\mu\text{-}OH)_2(\mu\text{-}OMe)(\mu_3\text{-}OMe)]^-$ (C. Jiang, Y.-S. Wen, L.-K. Liu, T.S.A. Hor and Y.K. Yan), 179

Pyrylium salts

Synthesis, structure and reactivity of acetylenic $Co_2(CO)_6$ pyrylium salts. Electronic influence of the $Co_2(CO)_6$ acetylenic fragment (B. Caro, F. Robin-Le Guen, M.-C. Sénéchal-Tocquer, V. Prat and J. Vaissermann), 87

Rhenium

Kohlenwasserstoffverbrückte Metallkomplexe, XXXVI. Nucleophile Addition von Carbonylmetallaten an koordinierte Ethen-, Benzol-, Cyclooctadienyl- und Cyclooctatrienyl-Liganden von kationischen Ruthenium-Komplexen (S. Hüffer, K. Polborn and W. Beck), 47

Ruthenium

Kohlenwasserstoffverbrückte Metallkomplexe, XXXVI. Nucleophile Addition von Carbonylmetallaten an koordinierte Ethen-, Benzol-, Cyclooctadienyl- und Cyclooctatrienyl-Liganden von kationischen Ruthenium-Komplexen (S. Hüffer, K. Polborn and W. Beck), 47

The preparation and reactions of $(\eta^3\text{-allyl})_2Ru[Pr^i_2P(CH_2)_nPPr^i_2]$ complexes (S. Holle, P.W. Jolly and J. Kuhnigk), 255

Selenium

Synthesis and spectroscopic characterization of $(CO)_6Fe_2\{\mu\text{-}EC(H){=}C(2\text{-}Th)E'\}$ (E,E' = S, Se, Te; Th = C_4H_3S). Structural characterization of $(CO)_6Fe_2\{\mu\text{-}SeC(H){=}C(2\text{-}Th)Se\}$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana, A.L. Rheingold, G.P.A. Yap and L.M. Liable-Sands), 135

Thermolysis of $Fe_3(CO)_9(\mu_3\text{-}Se)(\mu_3\text{-}E)$ (E = S, Te) with $Cp_2Mo_2(CO)_6$ and formation of new mixed-chalcogenide, mixed Fe–Mo carbonyl clusters. Crystal structures of $Cp_2Mo_2Fe_2(CO)_6(\mu_3\text{-}Se)_2(\mu_4\text{-}Te)$, $Cp_2Mo_2Fe_2(CO)_7(\mu_3\text{-}Se)(\mu_3\text{-}Te)$ and $Cp_2Mo_2Fe_2(CO)_6(\mu_3\text{-}S)(\mu_3\text{-}Se)(\mu_4\text{-}Se)$ (P. Mathur, S. Ghose, Md.M. Hossain, C.V.V. Satyanarayana and M.F. Mahon), 189

Silicon

Comparison of the reactivities of $(Me_3Si)_3CSiMe_2I$ and $(Pr^iMe_2Si)_3CSiMe_2I$ (A.I. Almansour), 83

On the reaction of $[Ph_2(OH)Si]_2O$ with t-Bu_2SnCl_2: synthesis and characterization of the first well defined polystannasiloxane $[(t\text{-}Bu_2SnO)(Ph_2SiO)_2]_n$ (J. Beckmann, K. Jurkschat, D. Schollmeyer and M. Schürmann), 229

Single crystal X-ray diffraction study

Methyl propiolate coupling at the $Fe(CO)_2(PEt_3)_2$ fragment (M. Jänicke, C. Gauss, A. Koller and H. Berke), 171

Steric hindrance

Comparison of the reactivities of $(Me_3Si)_3CSiMe_2I$ and $(Pr^iMe_2Si)_3CSiMe_2I$ (A.I. Almansour), 83

Styrene

Asymmetric hydroesterification of styrene by $PdCl_2$–$CuCl_2$-chiral phosphine catalyst systems (H. Zhou, J. Hou, J. Cheng, S. Lu, H. Fu and H. Wang), 227

Sulphur

Synthesis and spectroscopic characterization of $(CO)_6Fe_2\{\mu\text{-}EC(H){=}C(2\text{-}Th)E'\}$ (E,E' = S, Se, Te; Th = C_4H_3S). Structural

characterization of $(CO)_6Fe_2\{\mu$-SeC(H)=C(2-Th)Se$\}$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana, A.L. Rheingold, G.P.A. Yap and L.M. Liable-Sands), 135

Thermolysis of $Fe_3(CO)_9(\mu_3$-Se$)(\mu_3$-E) (E = S, Te) with $Cp_2Mo_2(CO)_6$ and formation of new mixed-chalcogenide, mixed Fe–Mo carbonyl clusters. Crystal structures of $Cp_2Mo_2Fe_2(CO)_6(\mu_3$-Se$)_2(\mu_4$-Te), $Cp_2Mo_2Fe_2(CO)_7(\mu_3$-Se$)(\mu_3$-Te) and $Cp_2Mo_2Fe_2(CO)_6(\mu_3$-S$)(\mu_3$-Se$)(\mu_4$-Se) (P. Mathur, S. Ghose, Md.M. Hossain, C.V.V. Satyanarayana and M.F. Mahon), 189

Synthesis

On the reaction of $[Ph_2(OH)Si]_2O$ with t-Bu_2SnCl_2: synthesis and characterization of the first well defined polystannasiloxane $[(t\text{-}Bu_2SnO)(Ph_2SiO)_2]_n$ (J. Beckmann, K. Jurkschat, D. Schollmeyer and M. Schürmann), 229

Synthesis of trinuclear mixed-metal clusters $Os_2Fe(CO)_{10}L_2$ (L = OH, I, CO). The crystal structures of $Os_2Fe(CO)_{10}(\mu$-OH$)_2$ and $Os_2Fe(CO)_{12} \cdot 0.41Os_3(CO)_{12}$ (V.A. Maksakov, V.P. Kirin, A.V. Virovets, N.V. Podberezskaya and P.P. Semyannikov), 57

Tellurium

Synthesis and spectroscopic characterization of $(CO)_6Fe_2\{\mu$-EC(H)=C(2-Th)E′$\}$ (E,E′ = S, Se, Te; Th = C_4H_3S). Structural characterization of $(CO)_6Fe_2\{\mu$-SeC(H)=C(2-Th)Se$\}$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana, A.L. Rheingold, G.P.A. Yap and L.M. Liable-Sands), 135

Thermolysis of $Fe_3(CO)_9(\mu_3$-Se$)(\mu_3$-E) (E = S, Te) with $Cp_2Mo_2(CO)_6$ and formation of new mixed-chalcogenide, mixed Fe–Mo carbonyl clusters. Crystal structures of $Cp_2Mo_2Fe_2(CO)_6(\mu_3$-Se$)_2(\mu_4$-Te), $Cp_2Mo_2Fe_2(CO)_7(\mu_3$-Se$)(\mu_3$-Te) and $Cp_2Mo_2Fe_2(CO)_6(\mu_3$-S$)(\mu_3$-Se$)(\mu_4$-Se) (P. Mathur, S. Ghose, Md.M. Hossain, C.V.V. Satyanarayana and M.F. Mahon), 189

Tellurium-oxide dimer

Isolation and X-ray structure determination of a new tellurane *Te*-oxide dimer $[12\text{-}Te\text{-}6(C4O2)]_2$ (λ^6-tellane) (S. Sato, T. Ueminami, E. Horn and N. Furukawa), 77

Thermal elimination

Osmium nitrosyl complexes with osmium–tin bonds. Crystal structure of $Os[Sn(p\text{-tolyl})_3](NO)(CO)_2(PPh_3)$ (A.M. Clark, C.E.F. Rickard, W.R. Roper and L.J. Wright), 111

Thiopheneacetylene

Synthesis and spectroscopic characterization of $(CO)_6Fe_2\{\mu$-EC(H)=C(2-Th)E′$\}$ (E,E′ = S, Se, Te; Th = C_4H_3S). Structural characterization of $(CO)_6Fe_2\{\mu$-SeC(H)=C(2-Th)Se$\}$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana, A.L. Rheingold, G.P.A. Yap and L.M. Liable-Sands), 135

Tin

On the reaction of $[Ph_2(OH)Si]_2O$ with t-Bu_2SnCl_2: synthesis and characterization of the first well defined polystannasiloxane $[(t\text{-}Bu_2SnO)(Ph_2SiO)_2]_n$ (J. Beckmann, K. Jurkschat, D. Schollmeyer and M. Schürmann), 229

Trimethylamine *N*-oxide

Amine-oxide-mediated reactions of $Re_2(CO)_{10}$ with phenol and aliphatic alcohols: The formation of $Re_3(CO)_{14}(\mu$-H) and a hydroxo-methoxo trirhenium aggregate $[Me_3NH]^+[Re_3(CO)_9(\mu$-OH$)_2(\mu$-OMe$)(\mu_3$-OMe$)]^-$ (C. Jiang, Y.-S. Wen, L.-K. Liu, T.S.A. Hor and Y.K. Yan), 179

Tungsten

Kohlenwasserstoffverbrückte Metallkomplexe, XXXV. Addition von *tert*. Butylethinyl–pentacarbonyl–wolframat an π-koordinierte Kohlenwasserstoffe von kationischen Komplexen (J. Milke, K. Sünkel and W. Beck), 39

Vinyliden bridged complexes

Kohlenwasserstoffverbrückte Metallkomplexe, XXXV. Addition von *tert*. Butylethinyl–pentacarbonyl–wolframat an π-koordinierte Kohlenwasserstoffe von kationischen Komplexen (J. Milke, K. Sünkel and W. Beck), 39

X-ray

Synthesis and spectroscopic characterization of $(CO)_6Fe_2\{\mu$-EC(H)=C(2-Th)E′$\}$ (E,E′ = S, Se, Te; Th = C_4H_3S). Structural characterization of $(CO)_6Fe_2\{\mu$-SeC(H)=C(2-Th)Se$\}$ (P. Mathur, A.K. Dash, Md.M. Hossain, C.V.V. Satyanarayana, A.L. Rheingold, G.P.A. Yap and L.M. Liable-Sands), 135

X-ray diffraction

On the reaction of $[Ph_2(OH)Si]_2O$ with t-Bu_2SnCl_2: synthesis and characterization of the first well defined polystannasiloxane $[(t\text{-}Bu_2SnO)(Ph_2SiO)_2]_n$ (J. Beckmann, K. Jurkschat, D. Schollmeyer and M. Schürmann), 229

X-ray structure

Synthesis of trinuclear mixed-metal clusters $Os_2Fe(CO)_{10}L_2$ (L = OH, I, CO). The crystal structures of $Os_2Fe(CO)_{10}(\mu$-OH$)_2$ and $Os_2Fe(CO)_{12} \cdot 0.41Os_3(CO)_{12}$ (V.A. Maksakov, V.P. Kirin, A.V. Virovets, N.V. Podberezskaya and P.P. Semyannikov), 57

X-ray structure analysis

Kooperative Wirkung in π-Ligand-verbrückten Zweikernkomplexen XX. Eine neue Strategie zur Synthese von heterodinuklearen Komplexen mit einem verbrückenden Bis(cyclopentadienyl)-Liganden [2] (J. Heck und J. Körnich), 153

Zirconium

Dinuclear cationic zirconium complexes with the fulvalene ligand. Synthesis and reactivity (T. Cuenca, M. Galakhov, G. Jiménez, E. Royo, P. Royo and M. Bochmann), 209